

Compassion Fatigue and Vicarious Trauma - Signs and Symptoms

Physical Signs and Symptoms

- Exhaustion
- Insomnia
- Headaches
- Increased susceptibility to illness
- Somatization and hypochondria

Behavioral Signs and Symptoms

- Increased use of alcohol and drugs
- Absenteeism
- Anger and Irritability
- Avoidance of clients
- Impaired ability to make decisions
- Problems in personal relationships
- Attrition
- Compromised care for clients
- The Silencing Response
- Depleted parenting

Psychological Signs and Symptoms

- Emotional exhaustion
- Distancing
- Negative self-image
- Depression
- Sadness, Loss of hope
- Anxiety
- Guilt
- Reduced ability to feel sympathy and empathy
- Cynicism
- Resentment
- Dread of working with certain clients
- Feeling professional helplessness
- Diminished sense of enjoyment/career
- Depersonalization/numbness
- Disruption of world view/ Heightened anxiety or irrational fears
- Inability to tolerate strong feelings
- Problems with intimacy
- Intrusive imagery – preoccupation with trauma
- Hypersensitivity to emotionally charged stimuli
- Insensitivity to emotional material
- Difficulty separating personal and professional lives
- Failure to nurture and develop non work related aspects of life

Sources: Saakvitne (1995), Figley (1995), Gentry, Baranowsky & Dunning (1997), Yassen (1995).

This sheet may be freely copied as long as (a) this box is left intact on the handout, (b) the author is credited, (c) no changes are made, and (d) it is not sold. Please be advised that compassion fatigue can lead to serious problems such as depression, anxiety and suicidal thoughts. The information contained on this sheet is not intended as a substitute for professional medical advice.

Copyright 2013 by Françoise Mathieu, www.compassionfatigue.ca